

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/GB05/000266

International filing date: 27 January 2005 (27.01.2005)

Document type: Certified copy of priority document

Document details: Country/Office: GB
Number: 0402138.2
Filing date: 30 January 2004 (30.01.2004)

Date of receipt at the International Bureau: 25 February 2005 (25.02.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



PUT/GB2005/000266.



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Stephen Hordley

Dated 11 February 2005




The
**Patent
Office**

1/77

02FEB04 EB69798-3 D02029
P01/7700 0.00-0402138.2 NONE

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office
Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference

ARG/NM/PB60707P

2. Patent application number

(The Patent Office will fill in his part)

0402138.2

30 JAN 2004

3. Full name, address and postcode of the or of each applicant (underline all surnames)

SmithKline Beecham Corporation
One Franklin Plaza, P.O. Box 7929, Philadelphia,
Pennsylvania 19101, United States of America

Patents ADP number (*if you know it*)

If the applicant is a corporate body, give the country/state of its incorporation

United States of America

5949417004

4. Title of the invention

Novel Compounds

5. Name of your agent (*if you have one*)

Corporate Intellectual Property

"Address for service" in the United Kingdom to which all correspondence should be sent
(including the postcode)

GlaxoSmithKline
Corporate Intellectual Property (CN9 25.1)
980 Great West Road
BRENTFORD
Middlesex TW8 9GS

Patents ADP number (*if you know it*)

8072555006

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and (*if you know it*) the or each application number

Country Priority application number Date of filing
(*if you know it*) (*day / month / year*)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application Date of filing
(*day / month / year*)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer yes if:*

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is named as an applicant, or
 - c) any named applicant is a corporate body
- See note (d)

9. Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form	0
Description	38
Claim(s)	5
Abstract	0
Drawings	0

8

10. If you are also filing any of the following, state how many against each item.

Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11. We request the grant of a patent on the basis of this application
Signature A R Gladwin Date 30-Jan-04

12. Name and daytime telephone number of person to contact in the United Kingdom
A R Gladwin 01438 762051

Warning

After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission unless an application has been filed at least six weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.


Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- For details of the fee and ways to pay please contact the Patent Office.

NOVEL COMPOUNDS

This invention relates to novel compounds and their use as pharmaceuticals, particularly as p38 kinase inhibitors, for the treatment of conditions or disease states mediated by p38 kinase activity or mediated by cytokines produced by the activity of p38 kinase.

We have now found a group of novel compounds that are inhibitors of p38 kinase. According to the invention there is provided a compound of formula (I):



(I)

wherein

A is a fused 5-membered heteroaryl ring optionally substituted by up to two substituents independently selected from C₁₋₆alkyl, -(CH₂)_k-C₃₋₇cycloalkyl, halogen, -CN, trifluoromethyl, -(CH₂)_kOR³, -(CH₂)_kCO₂R³, -(CH₂)_kNR³R⁴, -(CH₂)_kCONR³R⁴, -(CH₂)_kNHCOR³, -(CH₂)_kSO₂NR³R⁴, -(CH₂)_kNHSO₂R³, -(CH₂)_kSO₂(CH₂)_mR⁵, a 5- or 6-membered heterocyclyl ring containing nitrogen optionally substituted by C₁₋₂alkyl or -(CH₂)_kCO₂R³, and a 5-membered heteroaryl ring optionally substituted by C₁₋₂alkyl;

A is a fused 5-membered heteroaryl ring substituted by -BR⁶, and

A is optionally further substituted by one substituent selected from -OR⁷, halogen, trifluoromethyl, -CN, -CO₂R⁷ and C₁₋₆alkyl optionally substituted by hydroxy;

A is a fused 5-membered heteroaryl ring substituted by -(CH₂)_nheterocyclyl wherein the heterocyclyl is a 5- or 6-membered heterocyclic ring containing one or two heteroatoms independently selected from oxygen, sulfur and nitrogen optionally substituted by up to two substituents independently selected from oxo, C₁₋₆alkyl, -(CH₂)_pphenyl, -OR⁷, -(CH₂)_pCO₂R⁷, -NR⁷R⁸ and -CONR⁷R⁸, and

A is optionally further substituted by one substituent selected from -OR⁷, halogen, trifluoromethyl, -CN, -CO₂R⁷ and C₁₋₆alkyl optionally substituted by hydroxy; or

A is a fused 5-membered heteroaryl ring substituted by -(CH₂)_qaryl or -(CH₂)_qheteroaryl wherein the aryl or heteroaryl is optionally substituted by one or more substituents independently selected from oxo, C₁₋₆alkyl, halogen, -CN, trifluoromethyl, -OR⁹, -(CH₂)_rCO₂R¹⁰, -NR⁹R¹⁰, -(CH₂)_rCONR⁹R¹⁰, -NHCOR⁹, -SO₂NR⁹R¹⁰, -NHSO₂R⁹ and -S(O)_sR⁹, and

A is optionally further substituted by one substituent selected from $-OR^7$, halogen, trifluoromethyl, $-CN$, $-CO_2R^7$ and C_{1-6} alkyl optionally substituted by hydroxy;

R^1 is selected from methyl and chloro;

R^2 is selected from $-NH-CO-R^{11}$ and $-CO-NH-(CH_2)_t-R^{12}$;

5 R^3 is selected from hydrogen, C_{1-6} alkyl optionally substituted by up to two OH groups, $-(CH_2)_k-C_{3-7}$ cycloalkyl, $-(CH_2)_k$ phenyl optionally substituted by R^{13} and/or R^{14} and $-(CH_2)_k$ heteroaryl optionally substituted by R^{13} and/or R^{14} ,

R^4 is selected from hydrogen and C_{1-6} alkyl, or

10 R^3 and R^4 , together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and $N-R^{15}$;

R^5 is selected from C_{1-6} alkyl optionally substituted by up to three halogen atoms, C_{2-6} alkenyl optionally substituted by phenyl, C_{3-7} cycloalkyl, heteroaryl optionally substituted by up to three R^{13} and/or R^{14} groups, and phenyl optionally substituted by R^{13} and/or R^{14} ;

15 R^6 is a C_{3-6} alkyl group substituted by at least two substituents independently selected from $-OR^{16}$, $-NR^{16}R^{17}$, $-CO_2R^{16}$, $-CONR^{16}R^{17}$, $-NHCOR^{16}$ and $-NHSO_2R^{16}$;

R^7 and R^8 are each independently selected from hydrogen and C_{1-6} alkyl;

20 R^9 is selected from hydrogen, $-(CH_2)_u-C_{3-7}$ cycloalkyl, $-(CH_2)_u$ heterocyclyl, $-(CH_2)_u$ aryl, and C_{1-6} alkyl optionally substituted by up to two substituents independently selected from $-OR^{18}$ and $-NR^{18}R^{19}$,

R^{10} is selected from hydrogen and C_{1-6} alkyl, or

25 R^9 and R^{10} , together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and $N-R^{15}$;

R^{11} is selected from hydrogen, C_{1-6} alkyl, $-(CH_2)_t-C_{3-7}$ cycloalkyl, trifluoromethyl, $-(CH_2)_v$ heteroaryl optionally substituted by R^{20} and/or R^{21} , and $-(CH_2)_v$ phenyl optionally substituted by R^{20} and/or R^{21} ;

30 R^{12} is selected from hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, $-CONHR^{22}$, phenyl optionally substituted by R^{20} and/or R^{21} , and heteroaryl optionally substituted by R^{20} and/or R^{21} ;

R^{13} and R^{14} are each independently selected from halogen, $-CN$, trifluoromethyl, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, $-CONR^{22}R^{23}$, $-COR^{24}$, $-CO_2R^{24}$, and heteroaryl, or

35 R^{13} and R^{14} are linked to form a fused 5-membered heterocyclyl ring containing one heteroatom selected from oxygen, sulfur and $N-R^{15}$, or a fused heteroaryl ring;

R^{15} is selected from hydrogen and methyl;

R^{16} , R^{17} , R^{18} and R^{19} are each independently selected from hydrogen and C_{1-6} alkyl;

40 R^{20} is selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_t-C_{3-7}$ cycloalkyl, $-CONR^{22}R^{23}$, $-NHCOR^{23}$, halogen, $-CN$, $-(CH_2)_wNR^{25}R^{26}$, trifluoromethyl, phenyl optionally substituted by one or more R^{21} groups, and heteroaryl optionally substituted by one or more R^{21} groups;

R²¹ is selected from C₁₋₆alkyl, C₁₋₆alkoxy, halogen, trifluoromethyl, and -
(CH₂)_wNR²⁵R²⁶;

R²² and R²³ are each independently selected from hydrogen and C₁₋₆alkyl, or

R²² and R²³, together with the nitrogen atom to which they are bound, form a 5-
or 6-membered heterocyclic ring optionally containing one additional heteroatom selected
from oxygen, sulfur and N-R¹⁵, wherein the ring may be substituted by up to two C₁₋₆
alkyl groups;

R²⁴ is C₁₋₆alkyl;

R²⁵ is selected from hydrogen, C₁₋₆alkyl and -(CH₂)_t-C₃₋₇cycloalkyl optionally
substituted by C₁₋₆alkyl,

R²⁶ is selected from hydrogen and C₁₋₆alkyl, or

R²⁵ and R²⁶, together with the nitrogen atom to which they are bound, form a 5-
or 6-membered heterocyclic ring optionally containing one additional heteroatom selected
from oxygen, sulfur and N-R¹⁵;

R²⁷ is hydrogen or C₁₋₆alkyl;

B is selected from a bond, oxygen, NH and S(O)_x;

X and Y are each independently selected from hydrogen, methyl and halogen;

Z is selected from halogen, C₁₋₆alkyl and -OR²⁷;


k, m and w are each independently selected from 0, 1, 2 and 3;

n, q, r, s, t and x are each independently selected from 0, 1 and 2; and


u and v are each independently selected from 0 and 1;

or a pharmaceutically acceptable derivative thereof.

In one embodiment, A includes fused 5-membered heteroaryl rings containing up
to two heteroatoms independently selected from oxygen, nitrogen and sulfur. In another
embodiment, A includes fused 5-membered heteroaryl rings containing up to two
heteroatoms independently selected from oxygen and nitrogen. In a further embodiment,
A includes 5-membered heteroaryl rings containing two heteroatoms independently
selected from oxygen and nitrogen, for example rings containing a nitrogen atom and one
additional heteroatom selected from oxygen and nitrogen. Examples of suitable A groups
include fused isoxazolyl, pyrazolyl and pyrrolyl rings such as those shown below:



Representative examples of A groups include fused isoxazolyl and pyrazolyl rings
such as those shown below:



Ring A may be optionally substituted by substituents located on any position on the ring. Preferably, ring A is substituted by one substituent.

In one embodiment, A is optionally substituted by up to two substituents independently selected from C_{1-4} alkyl, in particular methyl; $-(CH_2)_k-C_{3-7}$ cycloalkyl, in particular $-(CH_2)_k$ -cyclopropyl; $-(CH_2)_kOR^3$; $-(CH_2)_kCO_2R^3$; $-(CH_2)_kNR^3R^4$; $-(CH_2)_kCONR^3R^4$; $-(CH_2)_kNHCOR^3$; $-(CH_2)_kSO_2(CH_2)_mR^5$; and a 5- or 6-membered heterocyclyl ring containing nitrogen, in particular 4-piperidinyl, optionally substituted by C_{1-2} alkyl or $-(CH_2)_kCO_2R^3$.


In another embodiment, A is substituted by $-BR^6$.

In another embodiment, A is substituted by $-(CH_2)_n$ heterocyclyl wherein the heterocyclyl is a 5- or 6-membered heterocyclic ring containing one or two heteroatoms independently selected from oxygen, sulfur and nitrogen optionally substituted by up to two substituents independently selected from oxo, C_{1-6} alkyl, $-(CH_2)_p$ phenyl, $-OR^7$, $-(CH_2)_pCO_2R^7$, $-NR^7R^8$ and $-CONR^7R^8$. Typically, the heterocyclyl is a 5- or 6-membered heterocyclic ring containing one or two heteroatoms independently selected from oxygen and nitrogen wherein the heterocyclyl is optionally substituted by up to two substituents located on any position on the ring. For example, when the heterocyclyl contains a sulfur atom, the sulfur atom may have up to two oxo substituents. In one embodiment, the heterocyclyl is substituted by $-(CH_2)_n$ phenyl.


In a further embodiment, A is substituted by $-(CH_2)_q$ aryl or $-(CH_2)_q$ heteroaryl wherein the aryl or heteroaryl is optionally substituted by one or more substituents independently selected from oxo, C_{1-6} alkyl, halogen, $-CN$, trifluoromethyl, $-OR^9$, $-(CH_2)_rCO_2R^{10}$, $-NR^9R^{10}$, $-(CH_2)_rCONR^9R^{10}$, $-NHCOR^9$, $-SO_2NR^9R^{10}$, $-NHSO_2R^9$ and $-S(O)_sR^9$. Typically, the $-(CH_2)_q$ aryl group is $-(CH_2)_q$ phenyl and the $-(CH_2)_q$ heteroaryl group is a group wherein the heteroaryl is a 5- or 6-membered heteroaryl ring containing up to two heteroatoms independently selected from oxygen and nitrogen. The $-(CH_2)_q$ aryl and $-(CH_2)_q$ heteroaryl groups are optionally substituted and the substituents may be located on any position on the aryl or heteroaryl. Preferably, the aryl is optionally substituted by one or two substituents independently selected from C_{1-6} alkyl, in particular methyl, halogen, $-CN$, trifluoromethyl, $-OR^9$, $-NR^9R^{10}$, $-(CH_2)_rCONR^9R^{10}$ and $-S(O)_sR^9$. Preferably, the heteroaryl is optionally substituted by one or two substituents independently selected from oxo and C_{1-6} alkyl, in particular methyl.

Representative examples include compounds of formula (I) wherein A is substituted by $-(CH_2)_q$ heteroaryl wherein the heteroaryl is a 5- or 6-membered heteroaryl ring containing up to two heteroatoms independently selected from oxygen and nitrogen and is optionally substituted by oxo.


5 In the compounds of formula (I), the Z group may be located on any position on the benzene ring as shown in formulae (IA), (IB) and (IC) below.



(IA)



(IB)



(IC)

10 In one embodiment, the compound of formula (I) is a compound of formula (IB) wherein R^1 , R^2 , A, X, Y and Z are as hereinbefore defined.

A representative example of R^1 is methyl.

A representative example of R^2 is $-CO-NH-(CH_2)_t-R^{12}$.

15 In one embodiment, R^3 selected from hydrogen; C_{1-6} alkyl optionally substituted by up to two OH groups, in particular methyl, ethyl, n-propyl, isopropyl, t-butyl or 2,2-dimethylpropyl optionally substituted by up to two OH groups; $-(CH_2)_k-C_{3-7}$ cycloalkyl, in particular $-(CH_2)_k$ -cyclopropyl; $-(CH_2)_k$ phenyl optionally substituted by R^{13} and/or R^{14} ; and $-(CH_2)_k$ heteroaryl, in particular thiazolyl, optionally substituted by R^{13} and/or R^{14} .

In one embodiment, R^4 is selected from hydrogen and C_{1-4} alkyl such as methyl.

20 Alternatively, R^3 and R^4 , together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom

selected from oxygen, sulfur and N-R¹⁵, in particular a pyrrolidinyl, piperidinyl, piperazinyl or 4-methylpiperazinyl, or morpholinyl ring.

In one embodiment, R⁵ is selected from C₁₋₆alkyl optionally substituted by up to three halogen atoms, in particular methyl, ethyl, n-propyl, isopropyl, n-butyl and n-hexyl optionally substituted by up to three halogen atoms; C₂₋₆alkenyl optionally substituted by phenyl, in particular ethenyl optionally substituted by phenyl; C₃₋₇cycloalkyl, in particular cyclopropyl; heteroaryl optionally substituted by R¹³ and/or R¹⁴, in particular a 5-membered heteroaryl ring containing at least one heteroatom selected from oxygen, nitrogen and sulfur such as furyl, thienyl, isoxazolyl, imidazolyl or pyrazolyl optionally substituted by up to three R¹³ and/or R¹⁴ groups; and phenyl optionally substituted by R¹³ and/or R¹⁴.

In one embodiment, R⁶ is a C₃₋₆alkyl group substituted by from two to four substituents, for example two substituents, independently selected from -OR¹⁶, -NR¹⁶R¹⁷ and -CO₂R¹⁶.

In one embodiment, R⁷ and R⁸ are independently selected from hydrogen and C₁₋₄alkyl.

In one embodiment, R⁹ is selected from hydrogen; -(CH₂)_u-C₃₋₇cycloalkyl, in particular -(CH₂)_u-cyclohexyl; -(CH₂)_uheterocyclyl, in particular wherein the heterocyclyl is a 5 or 6 membered heterocyclyl containing one heteroatom selected from oxygen, nitrogen and sulfur such a tetrahydrofuran or tetrahydropyran; and C₁₋₆alkyl, in particular C₁₋₄alkyl such as methyl, ethyl, or n-propyl, optionally substituted by up to two substituents independently selected from -OR¹⁸ and -NR¹⁸R¹⁹.

In one embodiment, R¹⁰ is hydrogen.

Alternatively, R⁹ and R¹⁰, together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N-R¹⁵, in particular morpholinyl.

In one embodiment, R¹¹ is a -(CH₂)_vheteroaryl optionally substituted by R²⁰ and/or R²¹.

In one embodiment, R¹² is selected from C₃₋₇cycloalkyl, phenyl optionally substituted by R²⁰ and/or R²¹, and heteroaryl optionally substituted by R²⁰ and/or R²¹. A representative example of R¹² is C₃₋₆cycloalkyl, in particular cyclopropyl.

In one embodiment, R¹³ and R¹⁴ are each independently selected from halogen, in particular chlorine or fluorine; -CN; trifluoromethyl; nitro; C₁₋₄alkyl, in particular methyl, ethyl, n-propyl, isopropyl or n-butyl; C₁₋₄alkoxy, in particular methoxy; -CONR²²R²³; -COR¹⁵; -CO₂R¹⁵; and heteroaryl, in particular a 5-membered heteroaryl ring containing up to two heteroatoms independently selected from nitrogen and oxygen, for example isoxazolyl.

Alternatively, R¹³ and R¹⁴ are linked to form a fused 5-membered heterocyclyl ring containing one heteroatom selected from oxygen, sulfur and N-R¹⁵.

In one embodiment, R¹⁶, R¹⁷, R¹⁸ and R¹⁹ are each independently selected from hydrogen and C₁₋₄alkyl.

In one embodiment, R²⁰ and R²¹ are each independently C₁₋₄alkoxy or -
(CH₂)_wNR²⁵R²⁶.

In one embodiment, R²² and R²³ are each independently hydrogen or C₁₋₄alkyl.

In one embodiment, R²⁴ is C₁₋₄alkyl.

5 In one embodiment, R²⁵ and R²⁶, together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally further containing one additional oxygen atom.

In one embodiment, R²⁷ is hydrogen or C₁₋₂alkyl.

In one embodiment, B is a bond.

10 In one embodiment, X and Y are each independently selected from hydrogen, chlorine and fluorine. Representative examples of X include hydrogen and fluorine. A representative example of Y is hydrogen.

In one embodiment, Z is halogen, in particular fluorine.

In one embodiment, k and m are independently selected from 0, 1 and 2.

15 In one embodiment, n and r are independently 1.

In one embodiment, q and u are independently selected from 0 and 1.

In one embodiment, s is 2.

A representative example of t is 0.

In one embodiment, v and w are independently 0.

20 It is to be understood that the present invention covers all combinations of particular and preferred groups described hereinabove. It is also to be understood that the present invention encompasses compounds of formula (I) in which a particular group or parameter, for example R³, R⁴, R⁵, R⁷, R⁸, R⁹, R¹⁰, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²¹, R²², R²³, R²⁴, R²⁵, R²⁶, k, m, p, r, s, t, u or w may occur more than once. In
25 such compounds it will be appreciated that each group or parameter is independently selected from the values listed.

Particular compounds according to the invention include those mentioned in the Examples. Specific examples which may be mentioned include:

N-cyclopropyl-3-[5-fluoro-3-(4-pyridinyl)-1*H*-indazol-6-yl]-4-methylbenzamide; and

30 *N*-cyclopropyl-3-fluoro-5-[5-fluoro-3-(4-pyridinyl)-1,2-benzisoxazol-6-yl]-4-methylbenzamide;

and pharmaceutically acceptable derivatives thereof.

As used herein, the term "pharmaceutically acceptable" means a compound which is suitable for pharmaceutical use. Salts and solvates of compounds of the
35 invention which are suitable for use in medicine are those wherein the counterion or associated solvent is pharmaceutically acceptable. However, salts and solvates having non-pharmaceutically acceptable counterions or associated solvents are within the scope of the present invention, for example, for use as intermediates in the preparation of other compounds of the invention and their pharmaceutically acceptable salts and
40 solvates.

As used herein, the term "pharmaceutically acceptable derivative", means any pharmaceutically acceptable salt, solvate or prodrug, e.g. ester, of a compound of the

invention, which upon administration to the recipient is capable of providing (directly or indirectly) a compound of the invention, or an active metabolite or residue thereof. Such derivatives are recognizable to those skilled in the art, without undue experimentation. Nevertheless, reference is made to the teaching of Burger's Medicinal Chemistry and Drug Discovery, 5th Edition, Vol 1: Principles and Practice, which is incorporated herein by reference to the extent of teaching such derivatives. Preferred pharmaceutically acceptable derivatives are salts, solvates, esters, carbamates and phosphate esters. Particularly preferred pharmaceutically acceptable derivatives are salts, solvates and esters. Most preferred pharmaceutically acceptable derivatives are salts and esters, in particular salts.

The compounds of the present invention may be in the form of and/or may be administered as a pharmaceutically acceptable salt. For a review on suitable salts see Berge *et al.*, J. Pharm. Sci., 1977, 66, 1-19.

Typically, a pharmaceutical acceptable salt may be readily prepared by using a desired acid or base as appropriate. The salt may precipitate from solution and be collected by filtration or may be recovered by evaporation of the solvent.

Salts of the compounds of the present invention may, for example, comprise acid addition salts resulting from reaction of an acid with a nitrogen atom present in a compound of formula (I). Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention. Suitable addition salts are formed from acids which form non-toxic salts and examples are acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, ethanesulfonate, formate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydrogen phosphate, hydroiodide, hydroxynaphthoate, iodide, isethionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, monopotassium maleate, mucate, napsylate, nitrate, N-methylglucamine, oxalate, oxaloacetate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, piruvate, polygalacturonate, saccharate, salicylate, stearate, subacetate, succinate, sulfate, tannate, tartrate, teoclate, tosylate, triethiodide, trifluoroacetate and valerate.

Pharmaceutically acceptable base salts include ammonium salts such as a trimethylammonium salt, alkali metal salts such as those of sodium and potassium, alkaline earth metal salts such as those of calcium and magnesium and salts with organic bases, including salts of primary, secondary and tertiary amines, such as isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexyl amine and N-methyl-D-glucamine.

Those skilled in the art of organic chemistry will appreciate that many organic compounds can form complexes with solvents in which they are reacted or from which they are precipitated or crystallized. These complexes are known as "solvates". As used herein, the term "solvate" refers to a complex of variable stoichiometry formed by a solute

(in this invention, a compound) for the purpose of the invention may not interfere with the biological activity of the solute. Examples of suitable solvents include water, methanol, ethanol and acetic acid. Preferably the solvent used is a pharmaceutically acceptable solvent. Examples of suitable pharmaceutically acceptable solvents include water, ethanol and acetic acid. Most preferably the solvent used is water. A complex with water is known as a "hydrate". Solvates of the compounds of the invention are within the scope of the invention.

As used herein, the term "prodrug" means a compound which is converted within the body, e.g. by hydrolysis in the blood, into its active form that has medical effects. Pharmaceutically acceptable prodrugs are described in T. Higuchi and V. Stella, Prodrugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series; Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987; and in D. Fleisher, S. Ramon and H. Barbra "Improved oral drug delivery: solubility limitations overcome by the use of prodrugs", Advanced Drug Delivery Reviews (1996) 19(2) 115-130, each of which are incorporated herein by reference.

Prodrugs are any covalently bonded carriers that release a compound of formula (I) *in vivo* when such prodrug is administered to a patient. Prodrugs are generally prepared by modifying functional groups in a way such that the modification is cleaved, either by routine manipulation or *in vivo*, yielding the parent compound. Prodrugs include, for example, compounds of this invention wherein hydroxy or amine groups are bonded to any group that, when administered to a patient, cleaves to form the hydroxy or amine groups. Thus, representative examples of prodrugs include (but are not limited to) acetate, formate and benzoate derivatives of alcohol and amine functional groups of the compounds of formula (I). Further, in the case of a carboxylic acid (-COOH), esters may be employed, such as methyl esters, ethyl esters, and the like. Esters may be active in their own right and/or be hydrolysable under *in vivo* conditions in the human body. Suitable pharmaceutically acceptable *in vivo* hydrolysable ester groups include those which break down readily in the human body to leave the parent acid or its salt.

As used herein, the term "alkyl" refers to straight or branched hydrocarbon chains containing the specified number of carbon atoms. For example, C₁₋₆alkyl means a straight or branched alkyl containing at least 1, and at most 6, carbon atoms. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, isobutyl, isopropyl and t-butyl. A C₁₋₄alkyl group is preferred, for example methyl, ethyl, isopropyl or t-butyl. The said alkyl groups may be optionally substituted with one or more fluorine atoms for example, trifluoromethyl.

As used herein, the term "alkenyl" refers to straight or branched hydrocarbon chains containing the specified number of carbon atoms and containing at least one double bond. For example, C₂₋₆alkenyl means a straight or branched alkenyl containing at least 2, and at most 6, carbon atoms and containing at least one double bond. Examples of "alkenyl" as used herein include, but are not limited to ethenyl, 2-propenyl, 3-butenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-methylbut-2-enyl, 3-hexenyl and 1,1-dimethylbut-2-enyl.

As used herein, the term "alkoxy" refers to a straight or branched chain alkoxy groups containing the specified number of carbon atoms. For example, C₁₋₆alkoxy means a straight or branched alkoxy containing at least 1, and at most 6, carbon atoms. Examples of "alkoxy" as used herein include, but are not limited to methoxy, ethoxy, propoxy, prop-2-oxy, butoxy, but-2-oxy, 2-methylprop-1-oxy, 2-methylprop-2-oxy, pentoxy, or hexyloxy. A C₁₋₄alkoxy group is preferred, for example methoxy or ethoxy.

As used herein, the term "cycloalkyl" refers to a non-aromatic hydrocarbon ring containing the specified number of carbon atoms which may optionally contain up to one double bond. For example, C₃₋₇cycloalkyl means a non-aromatic ring containing at least three, and at most seven, ring carbon atoms. Examples of "cycloalkyl" as used herein include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. A C₃₋₆cycloalkyl group is preferred, for example, cyclopropyl, cyclopentyl or cyclohexyl.

As used herein, the term "aryl" refers to an aromatic carbocyclic ring such as phenyl, biphenyl or naphthyl. Preferably the aryl is phenyl.

As used herein, the terms "heteroaryl ring" and "heteroaryl", unless otherwise defined, refer to a monocyclic 5- to 7-membered unsaturated hydrocarbon ring containing at least one heteroatom independently selected from oxygen, nitrogen and sulfur. Preferably, the heteroaryl ring has five or six ring atoms. Examples of heteroaryl rings include, but are not limited to, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, imidazolyl, pyrazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl and triazinyl. The said ring may be optionally substituted by one or more substituents independently selected from C₁₋₆alkyl and oxy.

As used herein, the terms "heterocyclic ring" or "heterocyclyl", unless otherwise defined refer to a monocyclic 3- to 7-membered saturated hydrocarbon ring containing at least one heteroatom independently selected from oxygen, nitrogen and sulfur. Preferably, the heterocyclyl ring has five or six ring atoms. Examples of heterocyclyl groups include, but are not limited to, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, piperidyl, piperazinyl, morpholino, tetrahydropyranyl, tetrahydrofuranyl, and thiomorpholino. The said ring may be optionally substituted by one or more substituents independently selected from C₁₋₆alkyl and oxy.

As used herein, the terms "halogen" or "halo" refer to the elements fluorine, chlorine, bromine and iodine. Preferred halogens are fluorine, chlorine and bromine. A particularly preferred halogen is fluorine or chlorine.

As used herein, the term "optionally" means that the subsequently described event(s) may or may not occur, and includes both event(s) which occur and events that do not occur.

As used herein, the term "substituted" refers to substitution with the named substituent or substituents, multiple degrees of substitution being allowed unless otherwise stated.

Certain compounds of formula (I) may exist in stereoisomeric forms (e.g. they may contain one or more asymmetric carbon atoms or may exhibit cis-trans isomerism). The


individual stereoisomers (enantiomers and diastereomers) and mixtures of these are included within the scope of the present invention. The present invention also covers the individual isomers of the compounds represented by formula (I) as mixtures with isomers thereof in which one or more chiral centres are inverted. Likewise, it is understood that compounds of formula (I) may exist in tautomeric forms other than that shown in the formula and these are also included within the scope of the present invention.

Separation of diastereoisomers or cis and trans isomers may be achieved by conventional techniques, e.g. by fractional crystallisation, chromatography or H.P.L.C. A stereoisomeric mixture of the agent may also be prepared from a corresponding optically pure intermediate or by resolution, such as H.P.L.C. of the corresponding racemate using a suitable chiral support or by fractional crystallisation of the diastereoisomeric salts formed by reaction of the corresponding racemate with a suitable optically active acid or base, as appropriate.

Furthermore, some of the crystalline forms of the compounds of structure (I) may exist as polymorphs, which are included in the present invention.


The compounds of this invention may be made by a variety of methods, including standard chemistry. Any previously defined variable will continue to have the previously defined meaning unless otherwise indicated. Illustrative general synthetic methods are set out below and then specific compounds of the invention are prepared in the working Examples.

A compound of formula (I) may be prepared by reacting a compound of formula (II)




(II)

in which A and Z are as hereinbefore defined and Hal is halogen, in particular bromine, with a compound of formula (IIIA) or (IIIB)



(IIIA)




(IIIB)

in which R¹, R², X and Y are as hereinbefore defined,

5 in the presence of a catalyst, for example tetrakis(triphenylphosphine)palladium.

A compound of formula (II) wherein A is a fused pyrazolyl ring may, for example, be prepared by reacting a compound of formula (IV)



(IV)

10

in which Z, Hal, $-(CH_2)_q$ aryl and $-(CH_2)_q$ heteroaryl are as hereinbefore defined and Hal¹ is halogen, in particular fluorine, with a protected hydrazine derivative of formula (V)

15




(V)

in which P is a protecting group such as Boc,

followed by cyclisation in the presence of a base such as DBU.

20 A compound of formula (II) wherein A is a fused isoxazolyl ring may, for example, be prepared by reacting a compound of formula (IV) as hereinbefore defined with hydroxylamine, followed by cyclisation in the presence of a base such as DBU.

A compound of formula (IIIA) may be prepared by, for example, reacting a compound of formula (VI)



(VI)

25


in which R^1 , R^2 , X and Y are as hereinbefore defined and Hal^2 is halogen, in particular iodine,

with bis(pinnacolato)diboron, [1,1'-bis(diphenylphosphino)ferrocene] dichloropalladium (II) complex ($PdCl_2(ppdf)$) and potassium acetate in a solvent such as DMF.

5 A compound of formula (IIIB) may be prepared by, for example, reacting a compound of formula (VI) as hereinbefore defined, with n-butyl lithium and triisopropyl borate in a solvent such as THF.

When R^2 is $-NH-CO-R^{11}$, a compound of formula (VI) may be prepared by reacting an amine of formula (VII)

10



(VII)

in which R^1 , X, Y and Hal^2 are as hereinbefore defined, with an acid compound of formula (VIII)

15



(VIII)


in which R^{11} is as hereinbefore defined, under amide forming conditions.

20

Suitable amide forming conditions are well known in the art and include adding a base such as DIPEA to a mixture of the amine of formula (VII), the acid of formula (VIII), and HATU in a solvent such as DMF.

Alternatively, when R^2 is $-CO-NH-(CH_2)_t-R^{12}$, a compound of formula (VI) may readily be prepared from a corresponding acid compound of formula (IX)

25




(IX)

in which R^1 , X, Y and Hal^2 are as hereinbefore defined,

30

by converting the acid to an activated form of the acid, for example the acid chloride, by treatment with, for example, thionyl chloride, and then reacting the activated acid thus formed with an amine compound of formula (X)




(X)

in which R^{12} is as hereinbefore defined,
under amide forming conditions.

5 Suitable amide forming conditions are well known in the art and include treating a solution of the acid of formula (IX), or the activated form thereof, in for example DMF, with an amine of formula (X) in the presence of a base such as triethylamine.


10 Alternatively, a further general method comprises final stage modification of one compound of formula (I) into another compound of formula (I). Suitable functional group transformations for converting one compound of formula (I) into another compound of formula (I) are well known in the art and are described in, for instance, *Comprehensive Heterocyclic Chemistry II*, eds. A. R. Katritzky, C. W. Rees and E. F. V. Scriven (Pergamon Press, 1996), *Comprehensive Organic Functional Group Transformations*, eds. A.R. Katritzky, O. Meth-Cohn and C.W. Rees (Elsevier Science Ltd., Oxford, 1995),
15 *Comprehensive Organic Chemistry*, eds. D. Barton and W.D. Ollis (Pergamon Press, Oxford, 1979), and *Comprehensive Organic Transformations*, R.C. Larock (VCH Publishers Inc., New York, 1989).

For example, one general method for preparing the compounds of formula (I) comprises the reaction set out in Scheme 1 below.




Scheme 1


For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 2 below.



For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 3 below.




For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 4 below.




Scheme 4

For example, another method for preparing the compounds of formula (I)
 5 comprises the reactions set out in Scheme 5 below.




Scheme 5

For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 6 below.




Scheme 6

For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 7 below.




Scheme 7

- For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 8 below.



Scheme 8

For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 9 below.




5

Scheme 9

For example, a further method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 10 below.

10



Scheme 10

Those skilled in the art will appreciate that in the preparation of the compounds of the invention it may be necessary and/or desirable to protect one or more sensitive groups in the molecule to prevent undesirable side reactions. Suitable protecting groups for use according to the present invention are well known to those skilled in the art and may be used in a conventional manner. See, for example, "Protective groups in organic synthesis" by T.W. Greene and P.G.M. Wuts (John Wiley & sons 1991) or "Protecting Groups" by P.J. Kocienski (Georg Thieme Verlag 1994). Examples of suitable amino protecting groups include acyl type protecting groups (e.g. formyl, trifluoroacetyl, acetyl), aromatic urethane type protecting groups (e.g. benzyloxycarbonyl (Cbz) and substituted Cbz), aliphatic urethane protecting groups (e.g. 9-fluorenylmethoxycarbonyl (Fmoc), t-butylloxycarbonyl (Boc), isopropylloxycarbonyl, cyclohexylloxycarbonyl) and alkyl type protecting groups (e.g. benzyl, trityl, chlorotriyl). Examples of suitable oxygen protecting groups may include for example alkyl silyl groups, such as trimethylsilyl or tert-butyl dimethylsilyl; alkyl ethers such as tetrahydropyranyl or tert-butyl; or esters such as acetate.

Whilst it is possible for the compounds of the present invention to be administered as the raw chemical, the compounds of formula (I) and their pharmaceutically acceptable derivatives are conveniently administered in the form of pharmaceutical compositions eg when the agent is in admixture with a suitable pharmaceutical excipient, diluent and/or

carrier selected with regard to the intended route of administration and standard pharmaceutical practice.

Thus, in another aspect of the invention, we provide a pharmaceutical composition comprising at least one compound of formula (I) or a pharmaceutically acceptable derivative thereof, in association with one or more pharmaceutically acceptable excipients, diluents and/or carriers. The excipient, diluent or carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

According to a further aspect, the invention provides a pharmaceutical composition comprising, as active ingredient, at least one compound of the invention or a pharmaceutically acceptable derivative thereof, in association one or more pharmaceutically acceptable excipients, diluents and/or carriers for use in therapy, and in particular in the treatment of human or animal subjects suffering from a condition susceptible to amelioration by an inhibitor of p38 kinase.

The present invention also provides a pharmaceutical composition comprising a therapeutically effective amount of the compounds of the present invention and a pharmaceutically acceptable excipient, diluent and/or carrier (including combinations thereof).

There is further provided by the present invention a process of preparing a pharmaceutical composition, which process comprises mixing at least one compound of the invention or a pharmaceutically acceptable derivative thereof, together with a pharmaceutically acceptable excipient, diluent and/or carrier.

The pharmaceutical compositions may be for human or animal usage in human and veterinary medicine and will typically comprise any one or more of a pharmaceutically acceptable excipient, diluent or carrier. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A. R. Gennaro edit. 1985). The choice of pharmaceutical excipient, diluent or carrier can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as – or in addition to – the excipient, diluent or carrier any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s) and solubilising agent(s).

Preservatives, stabilisers, dyes and even flavouring agents may be provided in the pharmaceutical composition. Examples of preservatives include sodium benzoate, sorbic acid and esters of p-hydroxybenzoic acid. Antioxidants and suspending agents may be also used.

For some embodiments, the agents of the present invention may also be used in combination with a cyclodextrin. Cyclodextrins are known to form inclusion and non-inclusion complexes with drug molecules. Formation of a drug-cyclodextrin complex may modify the solubility, dissolution rate, bioavailability and/or stability property of a drug molecule. Drug-cyclodextrin complexes are generally useful for most dosage forms and administration routes. As an alternative to direct complexation with the drug the

cyclodextrin may be used as an auxiliary additive, e. g. as a carrier, diluent or solubiliser. Alpha-, beta- and gamma-cyclodextrins are most commonly used and suitable examples are described in WO 91/11172, WO 94/02518 and WO 98/55148.

5 The compounds of the invention may be milled using known milling procedures such as wet milling to obtain a particle size appropriate for tablet formation and for other formulation types. Finely divided (nanoparticulate) preparations of the compounds of the invention may be prepared by processes known in the art, for example see WO 02/00196 (SmithKline Beecham).

10 There may be different composition/formulation requirements dependent on the different delivery systems. By way of example, the pharmaceutical composition of the present invention may be formulated to be delivered using a mini-pump or by a mucosal route, for example, as a nasal spray or aerosol for inhalation or ingestible solution, or parenterally in which the composition is formulated by an injectable form, for delivery, by, for example, an intravenous, intramuscular or subcutaneous route. Alternatively, the
15 formulation may be designed to be delivered by both routes.

Where the agent is to be delivered mucosally through the gastrointestinal mucosa, it should be able to remain stable during transit through the gastrointestinal tract; for example, it should be resistant to proteolytic degradation, stable at acid pH and resistant to the detergent effects of bile.

20 Where appropriate, the pharmaceutical compositions can be administered by inhalation, in the form of a suppository or pessary, topically in the form of a lotion, solution, cream, ointment or dusting powder, by use of a skin patch, orally in the form of tablets containing excipients such as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing
25 flavouring or colouring agents, or they can be injected parenterally, for example intravenously, intramuscularly or subcutaneously. For parenteral administration, the compositions may be best used in the form of a sterile aqueous solution which may contain other substances, for example enough salts or monosaccharides to make the solution isotonic with blood. For buccal or sublingual administration the compositions may
30 be administered in the form of tablets or lozenges which can be formulated in a conventional manner.

The routes for administration (delivery) include, but are not limited to, one or more of: oral (e. g. as a tablet, capsule, or as an ingestible solution), topical, mucosal (e. g. as a nasal spray or aerosol for inhalation), nasal, parenteral (e. g. by an injectable
35 form), gastrointestinal, intraspinal, intraperitoneal, intramuscular, intravenous, intrauterine, intraocular, intradermal, intracranial, intratracheal, intravaginal, intracerebroventricular, intracerebral, subcutaneous, ophthalmic (including intravitreal or intracameral), transdermal, rectal, buccal, epidural and sublingual. It is to be understood that not all of the compounds need be administered by the same route. Likewise, if the
40 composition comprises more than one active component, then those components may be administered by different routes.

The compounds of formula (I) and their pharmaceutically acceptable salts and solvates may be formulated for administration in any suitable manner. They may, for example, be formulated for topical administration or administration by inhalation or, more preferably, for oral, transdermal or parenteral administration. The pharmaceutical composition may be in a form such that it can effect controlled release of the compounds of formula (I) and their pharmaceutically acceptable derivatives. In a preferred embodiment, the agents of the present invention are delivered systemically such as orally, buccally or sublingually. A particularly preferred method of administration, and corresponding formulation, is oral administration.

For oral administration, the pharmaceutical composition may take the form of, and be administered as, for example, tablets (including sub-lingual tablets) and capsules (each including timed release and sustained release formulations), ovules, pills, powders, granules, elixirs, tinctures, emulsions, solutions, syrups or suspensions prepared by conventional means with acceptable excipients for immediate-, delayed-, modified-, sustained-, pulsed- or controlled-release applications.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. The tablets may also contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, dibasic calcium phosphate and glycine, disintegrants such as starch (preferably corn, potato or tapioca starch), sodium starch glycollate, croscarmellose sodium and certain complex silicates, and granulation binders such as polyvinylpyrrolidone, hydroxypropylmethylcellulose (HPMC), hydroxypropylcellulose (HPC), sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, stearic acid, glyceryl behenate and talc may be included.

Solid compositions of a similar type may also be employed as fillers in gelatin capsules. Preferred excipients in this regard include lactose, starch, a cellulose, milk sugar or high molecular weight polyethylene glycols. For aqueous suspensions and/or elixirs, the agent may be combined with various sweetening or flavouring agents, colouring matter or dyes, with emulsifying and/or suspending agents and with diluents such as water, ethanol, propylene glycol and glycerin, and combinations thereof.

Powders are prepared by comminuting the compound to a suitable fine size and mixing with a similarly comminuted pharmaceutical carrier such as an edible carbohydrate, as, for example, starch or mannitol. Flavoring, preservative, dispersing and coloring agent can also be present.

Capsules can be made by preparing a powder mixture as described above, and filling formed gelatin sheaths. Glidants and lubricants such as colloidal silica, talc, magnesium stearate, calcium stearate or solid polyethylene glycol can be added to the powder mixture before the filling operation. A disintegrating or solubilizing agent such as agar-agar, calcium carbonate or sodium carbonate can also be added to improve the availability of the medicament when the capsule is ingested.

Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

Tablets are formulated, for example, by preparing a powder mixture, granulating or slugging, adding a lubricant and disintegrant and pressing into tablets. A powder mixture is prepared by mixing the compound, suitably comminuted, with a diluent or base as described above, and optionally, with a binder such as carboxymethylcellulose, an aliginat, gelatin, or polyvinyl pyrrolidone, a solution retardant such as paraffin, a resorption accelerator such as a quaternary salt and/or an absorption agent such as bentonite, kaolin or dicalcium phosphate. The powder mixture can be granulated by wetting with a binder such as syrup, starch paste, acadia mucilage or solutions of cellulosic or polymeric materials and forcing through a screen. As an alternative to granulating, the powder mixture can be run through the tablet machine and the result is imperfectly formed slugs broken into granules. The granules can be lubricated to prevent sticking to the tablet forming dies by means of the addition of stearic acid, a stearate salt, talc or mineral oil. The lubricated mixture is then compressed into tablets. The compounds of the present invention can also be combined with free flowing inert carrier and compressed into tablets directly without going through the granulating or slugging steps. A clear or opaque protective coating consisting of a sealing coat of shellac, a coating of sugar or polymeric material and a polish coating of wax can be provided. Dyestuffs can be added to these coatings to distinguish different unit dosages.

Oral fluids such as solution, syrups and elixirs can be prepared in dosage unit form so that a given quantity contains a predetermined amount of the compound. Syrups can be prepared by dissolving the compound in a suitably flavored aqueous solution, while elixirs are prepared through the use of a non-toxic alcoholic vehicle. Suspensions can be formulated by dispersing the compound in a non-toxic vehicle. Solubilizers and emulsifiers such as ethoxylated isostearyl alcohols and polyoxy ethylene sorbitol ethers, preservatives, flavor additives such as peppermint oil or saccharin, and the like can also be added.

Where appropriate, dosage unit formulations for oral administration can be microencapsulated. The formulation can also be prepared to prolong or sustain the release as for example by coating or embedding particulate material in polymers, wax or the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

The compounds of the present invention can also be administered in the form of liposome emulsion delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

5 Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran
10 copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or
15 amphipathic block copolymers of hydrogels.

The present invention includes pharmaceutical compositions containing 0.1 to 99.5%, more particularly, 0.5 to 90% of a compound of the formula (I) in combination with a pharmaceutically acceptable carrier.

20 Likewise, the composition may also be administered in nasal, ophthalmic, otic, rectal, topical, intravenous (both bolus and infusion), intraperitoneal, intraarticular, subcutaneous or intramuscular, inhalation or insufflation form, all using forms well known to those of ordinary skill in the pharmaceutical arts.

For transdermal administration, the pharmaceutical composition may be given in the form of a transdermal patch, such as a transdermal iontophoretic patch.

25 If the compound of the present invention is administered parenterally, then examples of such administration include one or more of: intravenously, intraarterially, intraperitoneally, intrathecally, intraventricularly, intraurethrally, intrasternally, intracranially, intramuscularly or subcutaneously administering the agent; and/or by using infusion techniques. For parenteral administration, the pharmaceutical composition may
30 be given as an injection or a continuous infusion (e.g. intravenously, intravascularly or subcutaneously). The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. For administration by injection these may take the form of a unit dose presentation or as a multidose presentation preferably
35 with an added preservative. Alternatively for parenteral administration the active ingredient may be in powder form for reconstitution with a suitable vehicle. For parenteral administration, the compound is best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood. The aqueous solutions should be suitably buffered (preferably to a pH
40 of from 3 to 9), if necessary. The preparation of suitable parenteral formulations under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art.

The compositions of the present invention may be administered by direct injection.

The compounds of the invention may also be formulated as a depot preparation.

~~Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection.~~ Thus, for example, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Alternatively the composition may be formulated for topical application, for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

For application topically to the skin, the agent of the present invention can be formulated as a suitable ointment containing the active compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water.

Alternatively, it can be formulated as a suitable lotion or cream, suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, a polyethylene glycol, liquid paraffin, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

For administration by inhalation the compounds according to the invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, a hydrofluoroalkane such as tetrafluoroethane or heptafluoropropane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

Alternatively, the compound of the present invention can be administered in the form of a suppository or pessary, or it may be applied topically in the form of a gel, hydrogel, lotion, solution, cream, ointment or dusting powder.

The compounds of the present invention may also be administered by the pulmonary or rectal routes. They may also be administered by the ocular route. For ophthalmic use, the compounds can be formulated as micronised suspensions in isotonic, pH adjusted, sterile saline, or, preferably, as solutions in isotonic, pH adjusted, sterile

saline, optionally in combination with a preservative such as a benzylalkonium chloride. Alternatively, they may be formulated in an ointment such as petrolatum.

5 The pharmaceutical compositions generally are administered in an amount effective for treatment or prophylaxis of a specific condition or conditions. Initial dosing in humans is accompanied by clinical monitoring of symptoms, such symptoms for the selected condition. In general, the compositions are administered in an amount of active agent of at least about 100 $\mu\text{g/kg}$ body weight. In most cases they will be administered in one or more doses in an amount not in excess of about 20 mg/kg body weight per day. Preferably, in most cases, dose is from about 100 $\mu\text{g/kg}$ to about 5 mg/kg body weight, daily. For administration 10 particularly to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.1 mg/kg to 10 mg/kg and typically around 1 mg/kg . It will be appreciated that optimum dosage will be determined by standard methods for each treatment modality and indication, taking into account the indication, its severity, route of administration, complicating conditions and the like. The physician in any event will 15 determine the actual dosage which will be most suitable for an individual and will vary with the activity of the specific compound to be employed, the metabolic stability and length of action of that compound, age, weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, severity of the particular condition and response of the particular individual. The effectiveness of a selected actual dose can readily 20 be determined, for example, by measuring clinical symptoms or standard anti-inflammatory indicia after administration of the selected dose. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention. For conditions or disease states as are treated by the present invention, maintaining 25 consistent daily levels in a subject over an extended period of time, e.g., in a maintenance regime, can be particularly beneficial. For oral and parenteral administration to humans, the daily dosage level of the agent may be in single or divided doses.

In another aspect, the present invention provides a compound of formula (I) or a pharmaceutically acceptable derivative thereof, for use in therapy.

30 The compounds of the present invention are generally inhibitors of the serine/threonine kinase p38 and are therefore also inhibitors of cytokine production which is mediated by p38 kinase. Within the meaning of the term "inhibitors of the serine/threonine kinase p38" are included those compounds that interfere with the ability of p38 to transfer a phosphate group from ATP to a protein substrate according to the 35 assay described below.

It will be appreciated that the compounds of the invention may be selective for one or more of the isoforms of p38, for example p38 α , p38 β , p38 γ and/or p38 δ . In one embodiment, the compounds of the invention selectively inhibit the p38 α isoform. In another embodiment, the compounds of the invention selectively inhibit the p38 β isoform. 40 In a further embodiment, the compounds of the invention selectively inhibit the p38 α and p38 β isoforms. Assays for determining the selectivity of compounds for the p38 isoforms are described in, for example, WO 99/61426, WO 00/71535 and WO 02/46158.

It is known that p38 kinase activity can be elevated (locally or throughout the body), p38 kinase can be incorrectly temporally active or expressed, p38 kinase can be expressed or active in an inappropriate location, ~~p38 kinase can be constitutively expressed~~, or p38 kinase expression can be erratic; similarly, cytokine production mediated by p38 kinase activity can be occurring at inappropriate times, inappropriate locations, or it can occur at detrimentally high levels.

Accordingly, the present invention provides a method for the treatment of a condition or disease state mediated by p38 kinase activity, or mediated by cytokines produced by the activity of p38 kinase, in a subject which comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable derivative thereof. The compound may be administered as a single or polymorphic crystalline form or forms, an amorphous form, a single enantiomer, a racemic mixture, a single stereoisomer, a mixture of stereoisomers, a single diastereoisomer or a mixture of diastereoisomers.

The present invention also provides a method of inhibiting cytokine production which is mediated by p38 kinase activity in a subject, e.g. a human, which comprises administering to said subject in need of cytokine production inhibition a therapeutic, or cytokine-inhibiting, amount of a compound of the present invention. The compound may be administered as a single or polymorphic crystalline form or forms, an amorphous form, a single enantiomer, a racemic mixture, a single stereoisomer, a mixture of stereoisomers, a single diastereoisomer or a mixture of diastereoisomers.

The present invention treats these conditions by providing a therapeutically effective amount of a compound of this invention. By "therapeutically effective amount" is meant a symptom-alleviating or symptom-reducing amount, a cytokine-reducing amount, a cytokine-inhibiting amount, a kinase-regulating amount and/or a kinase-inhibiting amount of a compound. Such amounts can be readily determined by standard methods, such as by measuring cytokine levels or observing alleviation of clinical symptoms. For example, the clinician can monitor accepted measurement scores for anti-inflammatory treatments. It will be appreciated that reference to treatment includes acute treatment or prophylaxis as well as the alleviation of established symptoms.

The compounds of the present invention can be administered to any subject in need of inhibition or regulation of p38 kinase or in need of inhibition or regulation of p38 mediated cytokine production. In particular, the compounds may be administered to mammals. Such mammals can include, for example, horses, cows, sheep, pigs, mice, dogs, cats, primates such as chimpanzees, gorillas, rhesus monkeys, and, most preferably, humans.

Thus, the present invention provides methods of treating or reducing symptoms in a human or animal subject suffering from, for example, rheumatoid arthritis, osteoarthritis, asthma, psoriasis, eczema, allergic rhinitis, allergic conjunctivitis, adult respiratory distress syndrome, chronic pulmonary inflammation, chronic obstructive pulmonary disease, chronic heart failure, silicosis, endotoxemia, toxic shock syndrome, inflammatory bowel disease, tuberculosis, atherosclerosis, neurodegenerative disease, Alzheimer's disease,

Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, epilepsy, multiple sclerosis, aneurism, stroke, irritable bowel syndrome, muscle degeneration, bone resorption diseases, osteoporosis, diabetes, reperfusion injury, graft vs. host reaction, allograft rejections, sepsis, systemic cachexia, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome (AIDS), malaria, leprosy, infectious arthritis, leishmaniasis, Lyme disease, glomerulonephritis, gout, psoriatic arthritis, Reiter's syndrome, traumatic arthritis, rubella arthritis, Crohn's disease, ulcerative colitis, acute synovitis, gouty arthritis, spondylitis, and non articular inflammatory conditions, for example, herniated/ruptured/prolapsed intervertebral disk syndrome, bursitis, tendonitis, tenosynovitis, fibromyalgic syndrome and other inflammatory conditions associated with ligamentous sprain and regional musculoskeletal strain, pain, for example that associated with inflammation and/or trauma, osteopetrosis, restenosis, thrombosis, angiogenesis, cancer including breast cancer, colon cancer, lung cancer or prostatic cancer, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable derivative thereof.

A further aspect of the invention provides a method of treatment of a human or animal subject suffering from rheumatoid arthritis, asthma, psoriasis, chronic pulmonary inflammation, chronic obstructive pulmonary disease, chronic heart failure, systemic cachexia, glomerulonephritis, Crohn's disease, neurodegenerative disease, Alzheimer's disease, Parkinson's disease, epilepsy and cancer including breast cancer, colon cancer, lung cancer and prostatic cancer, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable derivative thereof.

A further aspect of the invention provides a method of treatment of a human or animal subject suffering from rheumatoid arthritis, asthma, psoriasis, chronic pulmonary inflammation, chronic obstructive pulmonary disease, chronic heart failure, systemic cachexia, glomerulonephritis, Crohn's disease and cancer including breast cancer, colon cancer, lung cancer and prostatic cancer, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable derivative thereof.

A further aspect of the invention provides a method of treatment of a human or animal subject suffering from rheumatoid arthritis, asthma, chronic pulmonary inflammation, chronic obstructive pulmonary disease, neurodegenerative disease, Alzheimer's disease, Parkinson's disease and epilepsy which comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable derivative thereof.

A further aspect of the invention provides a method of treatment of a human or animal subject suffering from any type of pain including chronic pain, rapid onset of analgesis, neuromuscular pain, headache, cancer pain, acute and chronic inflammatory pain associated with osteoarthritis and rheumatoid arthritis, post operative inflammatory pain, neuropathic pain, diabetic neuropathy, trigeminal neuralgia, post-hepatic neuralgia,

inflammatory neuropathies and migraine pain which comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable derivative thereof.

5 A further aspect of the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable derivative thereof, in the manufacture of a medicament for use in the treatment of a condition or disease state mediated by p38 kinase activity or mediated by cytokines produced by p38 kinase activity.

10 The compounds of formula (I) and their derivatives may be employed alone or in combination with other therapeutic agents for the treatment of the above-mentioned conditions. The invention thus provides, in a further aspect, a combination comprising a compound of the invention or a pharmaceutically acceptable derivative thereof together with a further therapeutic agent.

15 In particular, in rheumatoid arthritis therapy, combination with other chemotherapeutic or antibody agents is envisaged. Combination therapies according to the present invention thus comprise the administration of at least one compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof and at least one other pharmaceutically active agent. The compound(s) of formula (I) or pharmaceutically acceptable salt(s) or solvate(s) thereof and the other pharmaceutically active agent(s) may be administered together or separately and, when administered separately, this may
20 occur separately or sequentially in any order. The amounts of the compound(s) of formula (I) or pharmaceutically acceptable salt(s) or solvate(s) thereof and the other pharmaceutically active agent(s) and the relative timings of administration will be selected in order to achieve the desired combined therapeutic effect. Appropriate doses will be readily appreciated by those skilled in the art. It will be appreciated that the amount of a
25 compound of the invention required for treatment will vary with the nature of the condition being treated and the age and condition of the patient and will ultimately be at the discretion of the attendant physician or veterinarian. Examples of other pharmaceutically active agents which may be employed in combination with compounds of formula (I) and their salts and solvates for rheumatoid arthritis therapy include: immunosuppressants such as
30 as amtolmetin guacil, mizoribine and rimexolone; anti-TNF α agents such as etanercept, infliximab, diacerein; tyrosine kinase inhibitors such as leflunomide; kallikrein antagonists such as subreum; interleukin 11 agonists such as oprelvekin; interferon beta 1 agonists; hyaluronic acid agonists such as NRD-101 (Aventis); interleukin 1 receptor antagonists such as anakinra; CD8 antagonists such as amiprilose hydrochloride; beta amyloid
35 precursor protein antagonists such as reumacon; matrix metalloprotease inhibitors such as cipemastat and other disease modifying anti-rheumatic drugs (DMARDs) such as methotrexate, sulphasalazine, cyclosporin A, hydroxychloroquine, auranofin, aurothioglucose, gold sodium thiomalate and penicillamine.

40 The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical formulations comprising a combination as defined above together with a pharmaceutically acceptable carrier or excipient comprise a further aspect of the invention.

The individual components of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations by any convenient route.

- 5 When administration is sequential, either the compound of the invention or the second therapeutic agent may be administered first. When administration is simultaneous, the combination may be administered either in the same or different pharmaceutical composition.


- 10 When combined in the same formulation it will be appreciated that the two compounds must be stable and compatible with each other and the other components of the formulation. When formulated separately they may be provided in any convenient formulation, conveniently in such manner as are known for such compounds in the art.

EXAMPLES

The following Examples are illustrative embodiments of the invention, not limiting the scope of the invention in any way. Reagents are commercially available or are prepared according to procedures in the literature.

LCMS was conducted on a column (3.3cm x 4.6mm ID, 3um ABZ+PLUS), at a Flow Rate of 3ml/min, Injection Volume of 5µl, at room temperature and UV Detection Range at 215 to 330nm.


Intermediate 1: Z/E 1-(4-Bromo-2,5-difluorophenyl)-1-(4-pyridinyl)methanimine



A stirred solution of 1,4-dibromo-2,5-difluorobenzene (2.72g) in anhydrous diethylether (50ml) under nitrogen was cooled to -78°C in a dry-ice/acetone bath then treated with n-butyllithium (1.6M in hexanes, 7.2ml). The reaction mixture was stirred for 20min and a solution of 4-cyanopyridine (1.4g) in diethylether (60ml) was added dropwise. The resulting mixture was stirred for a further 20min then treated with saturated aqueous ammonium chloride. The ether layer was separated, washed with brine, dried (sodium sulfate) and concentrated under vacuum. The crude product was purified on an SPE cartridge (silica) using a cyclohexane/ethyl acetate gradient as eluant to give the title compound as a colourless gum (1.50g).

NMR: [δ H CDCl₃] 10.35 (1H, s, minor), 10.35 (1H, s, major) 8.74(2H, d, J=6Hz, major and minor), 7.58 (2H, d, J=6Hz, major), 7.48-7.42(1H, m,), 7.40-7.03 (4H, m, major and minor).


Intermediate 2: 1-(4-Bromo-2,5-difluorophenyl)-1-(4-pyridinyl)methananone



A mixture of 1-(4-bromo-2,5-difluorophenyl)-1-(4-pyridinyl)methanimine (Intermediate 1, 1.30g) in methanol (20ml) was treated with 0.5M hydrochloric acid (1.5ml). The reaction mixture was left at room temperature for 18h then applied to an acidic ion exchange cartridge (SCX). Elution with methanol followed by 10% aqueous ammonia in methanol gave the title compound as a white solid (1.20g).

LC-MS: Rt 2.96, MH^+ 298/300.


Intermediate 3: 1,1-Dimethylethyl (2Z)-2-[(4-bromo-2,5-difluorophenyl)(4-pyridinyl)methylidene]hydrazinecarboxylate



A solution of 1-(4-bromo-2,5-difluorophenyl)-1-(4-pyridinyl)methanone (Intermediate 2, 900mg) and t-butylcarbazate (400mg) in methanol (6ml) and acetic acid (0.5ml) was stirred at reflux for 12h. The mixture was partitioned between water and ethyl acetate and the organic layer was washed with water and brine, dried using a hydrophobic filter tube and concentrated under vacuum. The residue was purified on an SPE cartridge (silica, 5g) eluting with cyclohexane/ethyl acetate (9:1 to 6:4) to give the title compound as a yellow gel (880mg).

LC-MS: Rt 3.1/3.26, MH^+ 412/414.


Intermediate 4: 6-Bromo-5-fluoro-3-(4-pyridinyl)-1H-indazole



A solution of 1,1-dimethylethyl (2Z)-2-[(4-bromo-2,5-difluorophenyl)(4-pyridinyl)methylidene]hydrazinecarboxylate (Intermediate 3, 250mg) and DBU (95μl) in THF (3ml) was stirred at 150°C in a microwave oven. After 30min the mixture was partitioned between water and ethyl acetate and the organic layer was washed with water and brine, dried using a hydrophobic filter tube and concentrated under vacuum to give the title compound as an orange solid (150mg).

LC-MS: Rt 2.7min, MH^+ 292/294.


Intermediate 5: Z/E (4-Bromo-2,5-difluorophenyl)(4-pyridinyl)methanone oxime



5 A solution of Z/E 1-(4-bromo-2,5-difluorophenyl)-1-(4-pyridinyl)methanimine
 (Intermediate 1, 150mg) in ethanol (2ml) was treated with a solution of hydroxylamine
 hydrochloride (44mg) and sodium acetate (50mg) in water (0.4ml) then stirred at reflux for
 2h. The ethanol was removed under vacuum and the residue was partitioned between
 water and ethyl acetate. The aqueous layer was re-extracted with ethyl acetate and the
 combined organic extracts were washed with brine, dried (sodium sulfate) and
 10 concentrated under vacuum. The residue was purified on an SPE cartridge (silica, 2g)
 eluting with cyclohexane/ethyl acetate (4:1 to 1:1) to give the title compound as a white
 foam (130mg).

LC-MS: 2.83min, 2.90min, MH+ 313, 315.


15 **Intermediate 6: 6-Bromo-5-fluoro-3-(4-pyridinyl)-1,2-benzisoxazole**



20 A solution of Z/E (4-bromo-2,5-difluorophenyl)(4-pyridinyl)methanone oxime
 (Intermediate 5, 124mg) and 1,8-diazabicyclo[5.4.0]undec-7-ene (40mg) in
 tetrahydrofuran (1.5ml) was heated at 150°C in a microwave oven for 30min. The solvent
 was removed under vacuum and the residue was triturated with a small quantity of
 methanol to give the title compound as a cream coloured solid (32mg).

LC-MS: Rt 3.21min, MH+ 293, 295.


25 **Example 1: N-Cyclopropyl-3-[5-fluoro-3-(4-pyridinyl)-1H-indazol-6-yl]-4-methylbenzamide**



A suspension of 6-bromo-5-fluoro-3-(4-pyridinyl)-1*H*-indazole (Intermediate 4, 20mg), *N*-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (21mg), tetrakis(triphenylphosphine)palladium(0) (2mg) and aqueous sodium hydrogen carbonate (1M, 1ml) in isopropanol (2ml) was stirred in a microwave oven at 150°C for 15min. The mixture was partitioned between water and ethyl acetate and the organic layer was washed with water and brine, dried using a hydrophobic filter tube and concentrated under vacuum. The residue was purified on an SPE cartridge (silica, 5g) eluting with cyclohexane:ethyl acetate (1:1 to 1:4) to give the title compound as a white solid (9.72mg).

LC-MS: Rt 2.64, MH⁺ 387.

Example 2: *N*-Cyclopropyl-3-[5-fluoro-3-(1-oxido-4-pyridinyl)-1*H*-indazol-6-yl]-4-methylbenzamide



A solution of *N*-cyclopropyl-3-[5-fluoro-3-(4-pyridinyl)-1*H*-indazol-6-yl]-4-methylbenzamide (Example 1, 9.3mg) and *m*-CPBA (20mg) in chloroform (2ml) was stirred under nitrogen at 60°C for 1h. The mixture was applied to an Isolute amino cartridge (1g) and eluted with chloroform/methanol (100:0 and 90:10) to give the title compound as a yellow solid (5.5mg).

LC-MS: Rt 2.58, MH⁺ 403.

Example 3: *N*-Cyclopropyl-3-fluoro-5-[5-fluoro-3-(4-pyridinyl)-1,2-benzisoxazol-6-yl]-4-methylbenzamide

	Boc	t-Butoxycarbonyl
	^t BuONO	t-Butyl nitrite
	CDI	Carbonyldiimidazole
	DBU	1,8-Diazabicyclo[5.4.0]undec-7-ene
5	DIPEA	N,N-Diisopropylethylamine
	DMF	Dimethylformamide
	Et ₂ O	Diethyl ether
	EtOH	Ethanol
	h	Hours
10	Hal	Halogen
	HATU	O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
	HOBt	1-Hydroxybenzotriazole hydrate
	KOAc	Potassium Acetate
15	m-CPBA	3-Chloroperbenzoic acid
	MeCN	Acetonitrile
	MeOH	Methanol
	min	Minutes
	Ms	Mesyl
20	NaO ^t Bu	Sodium tert-butoxide
	NaOEt	Sodium Ethoxide
	NCS	N-Chlorosuccinimide
	PdCl ₂ (dppf)	[1,1'-bis(Diphenylphosphino)ferrocene]dichloropalladium (II) complex with dichloromethane (1:1)
25	R _t	Retention Time
	SPE	Solid phase extraction
	THF	Tetrahydrofuran

BIOLOGICAL EXAMPLES

30

The activity of compounds of formula (I) as p38 inhibitors may be determined by the following *in vitro* assays:

Fluorescence anisotropy kinase binding assay

35


The kinase enzyme, fluorescent ligand and a variable concentration of test compound are incubated together to reach thermodynamic equilibrium under conditions such that in the absence of test compound the fluorescent ligand is significantly (>50%) enzyme bound and in the presence of a sufficient concentration (>10x K_i) of a potent inhibitor the anisotropy of the unbound fluorescent ligand is measurably different from the bound value.

40

The concentration of kinase enzyme should preferably be $\geq 1 \times K_f$. The concentration of fluorescent ligand required will depend on the instrumentation used, and

CLAIMS

1. A compound of formula (I):



5

(I)

wherein

A is a fused 5-membered heteroaryl ring optionally substituted by up to two substituents independently selected from C_{1-6} alkyl, $-(CH_2)_k-C_{3-7}$ cycloalkyl, halogen, -CN, trifluoromethyl, $-(CH_2)_kOR^3$, $-(CH_2)_kCO_2R^3$, $-(CH_2)_kNR^3R^4$, $-(CH_2)_kCONR^3R^4$, $-(CH_2)_kNHCOR^3$, $-(CH_2)_kSO_2NR^3R^4$, $-(CH_2)_kNHSO_2R^3$, $-(CH_2)_kSO_2(CH_2)_mR^5$, a 5- or 6-membered heterocyclyl ring containing nitrogen optionally substituted by C_{1-2} alkyl or $-(CH_2)_kCO_2R^3$, and a 5-membered heteroaryl ring optionally substituted by C_{1-2} alkyl;

A is a fused 5-membered heteroaryl ring substituted by $-BR^6$, and

A is optionally further substituted by one substituent selected from $-OR^7$, halogen, trifluoromethyl, -CN, $-CO_2R^7$ and C_{1-6} alkyl optionally substituted by hydroxy;

A is a fused 5-membered heteroaryl ring substituted by $-(CH_2)_n$ heterocyclyl wherein the heterocyclyl is a 5- or 6-membered heterocyclic ring containing one or two heteroatoms independently selected from oxygen, sulfur and nitrogen optionally substituted by up to two substituents independently selected from oxo, C_{1-6} alkyl, $-(CH_2)_p$ phenyl, $-OR^7$, $-(CH_2)_pCO_2R^7$, $-NR^7R^8$ and $-CONR^7R^8$, and

A is optionally further substituted by one substituent selected from $-OR^7$, halogen, trifluoromethyl, -CN, $-CO_2R^7$ and C_{1-6} alkyl optionally substituted by hydroxy; or

A is a fused 5-membered heteroaryl ring substituted by $-(CH_2)_q$ aryl or $-(CH_2)_q$ heteroaryl wherein the aryl or heteroaryl is optionally substituted by one or more substituents independently selected from oxo, C_{1-6} alkyl, halogen, -CN, trifluoromethyl, $-OR^9$, $-(CH_2)_rCO_2R^{10}$, $-NR^9R^{10}$, $-(CH_2)_rCONR^9R^{10}$, $-NHCOR^9$, $-SO_2NR^9R^{10}$, $-NHSO_2R^9$ and $-S(O)_sR^9$, and

A is optionally further substituted by one substituent selected from $-OR^7$, halogen, trifluoromethyl, -CN, $-CO_2R^7$ and C_{1-6} alkyl optionally substituted by hydroxy;

R^1 is selected from methyl and chloro;

R^2 is selected from $-NH-CO-R^{11}$ and $-CO-NH-(CH_2)_t-R^{12}$;

R^3 is selected from hydrogen, C_{1-6} alkyl optionally substituted by up to two OH groups, $-(CH_2)_k-C_{3-7}$ cycloalkyl, $-(CH_2)_k$ phenyl optionally substituted by R^{13} and/or R^{14} and $-(CH_2)_k$ heteroaryl optionally substituted by R^{13} and/or R^{14} ,

R^4 is selected from hydrogen and C_{1-6} alkyl, or

5 R^3 and R^4 , together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N- R^{15} ;

10 R^5 is selected from C_{1-6} alkyl optionally substituted by up to three halogen atoms, C_{2-6} alkenyl optionally substituted by phenyl, C_{3-7} cycloalkyl, heteroaryl optionally substituted by up to three R^{13} and/or R^{14} groups, and phenyl optionally substituted by R^{13} and/or R^{14} ;

R^6 is a C_{3-6} alkyl group substituted by at least two substituents independently selected from $-OR^{16}$, $-NR^{16}R^{17}$, $-CO_2R^{16}$, $-CONR^{16}R^{17}$, $-NHCOR^{16}$ and $-NHOSO_2R^{16}$;

R^7 and R^8 are each independently selected from hydrogen and C_{1-6} alkyl;

15 R^9 is selected from hydrogen, $-(CH_2)_u-C_{3-7}$ cycloalkyl, $-(CH_2)_u$ heterocyclyl, $-(CH_2)_u$ aryl, and C_{1-6} alkyl optionally substituted by up to two substituents independently selected from $-OR^{18}$ and $-NR^{18}R^{19}$,

R^{10} is selected from hydrogen and C_{1-6} alkyl, or

20 R^9 and R^{10} , together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N- R^{15} ;

R^{11} is selected from hydrogen, C_{1-6} alkyl, $-(CH_2)_t-C_{3-7}$ cycloalkyl, trifluoromethyl, $-(CH_2)_v$ heteroaryl optionally substituted by R^{20} and/or R^{21} , and $-(CH_2)_v$ phenyl optionally substituted by R^{20} and/or R^{21} ;

25 R^{12} is selected from hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, $-CONHR^{22}$, phenyl optionally substituted by R^{20} and/or R^{21} , and heteroaryl optionally substituted by R^{20} and/or R^{21} ;

R^{13} and R^{14} are each independently selected from halogen, $-CN$, trifluoromethyl, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, $-CONR^{22}R^{23}$, $-COR^{24}$, $-CO_2R^{24}$, and heteroaryl, or

30 R^{13} and R^{14} are linked to form a fused 5-membered heterocyclyl ring containing one heteroatom selected from oxygen, sulfur and N- R^{15} , or a fused heteroaryl ring;

R^{15} is selected from hydrogen and methyl;

R^{16} , R^{17} , R^{18} and R^{19} are each independently selected from hydrogen and C_{1-6} alkyl;

35 R^{20} is selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_t-C_{3-7}$ cycloalkyl, $-CONR^{22}R^{23}$, $-NHCOR^{23}$, halogen, $-CN$, $-(CH_2)_wNR^{25}R^{26}$, trifluoromethyl, phenyl optionally substituted by one or more R^{21} groups, and heteroaryl optionally substituted by one or more R^{21} groups;

40 R^{21} is selected from C_{1-6} alkyl, C_{1-6} alkoxy, halogen, trifluoromethyl, and $-(CH_2)_wNR^{25}R^{26}$;

R^{22} and R^{23} are each independently selected from hydrogen and C_{1-6} alkyl, or

R²² and R²³, together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N-R¹⁵, wherein the ring may be substituted by up to two C₁₋₆alkyl groups;

5 R²⁴ is C₁₋₆alkyl;

R²⁵ is selected from hydrogen, C₁₋₆alkyl and -(CH₂)_t-C₃₋₇cycloalkyl optionally substituted by C₁₋₆alkyl,

R²⁶ is selected from hydrogen and C₁₋₆alkyl, or

10 R²⁵ and R²⁶, together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N-R¹⁵;

R²⁷ is hydrogen or C₁₋₆alkyl;

B is selected from a bond, oxygen, NH and S(O)_x;

X and Y are each independently selected from hydrogen, methyl and halogen;

15 Z is selected from halogen, C₁₋₆alkyl and -OR²⁷;

k, m and w are each independently selected from 0, 1, 2 and 3;

n, q, r, s, t and x are each independently selected from 0, 1 and 2; and

u and v are each independently selected from 0 and 1;

or a pharmaceutically acceptable derivative thereof.

20

2. A compound according to claim 1 wherein A is a fused 5-membered heteroaryl ring containing up to two heteroatoms independently selected from oxygen and nitrogen.

25 3. A compound according to claim 1 or claim 2 wherein A is substituted by -(CH₂)_qaryl or -(CH₂)_qheteroaryl wherein the aryl or heteroaryl is optionally substituted by one or more substituents independently selected from oxo, C₁₋₆alkyl, halogen, -CN, trifluoromethyl, -OR⁹, -(CH₂)_rCO₂R¹⁰, -NR⁹R¹⁰, -(CH₂)_rCONR⁹R¹⁰, -NHCOR⁹, -SO₂NR⁹R¹⁰, -NH₂SO₂R⁹ and -S(O)_sR⁹.

30 4. A compound according to anyone of the preceding claims wherein R¹ is methyl.

5. A compound according to any one of the preceding claims wherein R² is -CO-NH-(CH₂)_t-R¹².

35 6. A compound according to any one of the preceding claims wherein X is hydrogen or fluorine.

7. A compound according to claim 1 substantially as hereinbefore defined with reference to any one of Examples 1 to 4, or a pharmaceutically acceptable derivative thereof.

40

8. A compound according to claim 1 selected from:

N-cyclopropyl-3-[5-fluoro-3-(4-pyridinyl)-1*H*-indazol-6-yl]-4-methylbenzamide; and
N-cyclopropyl-3-fluoro-5-[5-fluoro-3-(4-pyridinyl)-1,2-benzisoxazol-6-yl]-4-methylbenzamide;
 or a pharmaceutically acceptable derivative thereof.

5

9. A pharmaceutical composition comprising at least one compound as claimed in any one of claims 1 to 8, or a pharmaceutically acceptable derivative thereof, in association with one or more pharmaceutically acceptable excipients, diluents and/or carriers.

10

10. A compound according to any one of claims 1 to 8, or a pharmaceutically acceptable derivative thereof, for use in therapy.

15

11. A method for treating a condition or disease state mediated by p38 kinase activity or mediated by cytokines produced by the activity of p38 kinase comprising administering to a patient in need thereof a compound as claimed in any one of claims 1 to 8, or a pharmaceutically acceptable derivative thereof.


20

12. Use of a compound as claimed in any one of claims 1 to 8, or a pharmaceutically acceptable derivative thereof, in the manufacture of a medicament for use in the treatment of a condition or disease state mediated by p38 kinase activity or mediated by cytokines produced by the activity of p38 kinase.

25

13. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 8, or a pharmaceutically acceptable derivative thereof, which comprises


(a) reacting a compound of formula (II)




(II)

30

in which A is defined in claim 1 and Hal is halogen,
 with a compound of formula (IIIA) or (IIIB)



(IIIA)



(IIIB)

5

in which R^1 , R^2 , X and Y are as defined in claim 1,
in the presence of a catalyst, or

10

(b) final stage modification of one compound of formula (I) as defined in claim 1 to
give another compound of formula (I) as defined in claim 1.



1